

# **Mouse Anti Human CD14 Purified**

## PRODUCT INFORMATION

CLONE: HI221

**ISOTYPE**: Mouse IgM **WS.No**.: VI MA52

**CATALOG#**: A6281/ A6281-bulk

**CONTENTS:** Purified antibody buffered in 10mM PBS (PH 7.0) with 0.1% NaN<sub>3</sub>.

### **DESCRIPTION**

CD14 McAb recognizes a 53-55KD glycosylphosphatidy-linositol (GPI)-anchored single chain glycoprotein, which is found on cells of myelomonocytic lineage including monocytes(strongly), granulocytes(weakly) and also on most tissue macrophages. CD14 antigen is also present on reticular dendritic cells, some langerhans cells, and B lymphocytes (weakly), but is absent from T lymphocytes, NK cells, erythrocytes and platelets. CD14 functions as a high affinity receptor for the complex of lipopolysaccharide (LPS-endotoxin) and LPS-binding protein (LBP), and is the major receptor involved in the lethal response to both endotoxin and Gram-negative bacteria. In addition to membrane CD14, soluble form of CD14 can be detected in serum and tissue culture supernatants of cells transfected with CD14.

#### **PREPARATION**

The monoclonal antibody is purified from ascites by hydroxyapatite chromatography.

## **USAGE**

The purified reagent is effective for indirect immunofluorescence staining of human cells for flow cytometric analysis.

#### **STORAGE**

Store at 4℃. For long-term storage aliquot and store at -20℃. Avoid freeze and thaw cycles.

#### **REFERENCES**

Tadamitsu. K., K.Hitoshi, A.E.G.Kr.van dem Borne, et al., eds. 1997. Leucocyte Typing W: White Cell Differentiation Antigens. P: 940, 963-968. Garland Publishing, Inc., New York.